

FARNAM COMPANIES, INC.

301 WEST OSBORN ROAD, P.O. BOX 34820, PHOENIX, AZ, 85013-3928

Telephone: 602-285-1660

Toll-Free: 800-234-2269

Fax: 602-207-2183

Website: www.farnam.com

ADAMS™ PLUS FLEA & TICK COLLAR FOR SMALL DOGS

Farnam

SMALL DOGS

Up to 5 Months Protection

KILLS:

fleas

ticks

flea eggs

flea larvae

ACTIVE INGREDIENTS:

Propoxur (CAS #114-26-1)	10.0%
(S)-Methoprene (CAS #65733-16-6)	2.1%
OTHER INGREDIENTS:	87.9%
TOTAL:	100.0%

DO NOT ALLOW CHILDREN TO PLAY WITH THIS COLLAR

CAUTION

READ ENTIRE LABEL BEFORE EACH USE.

USE ONLY ON DOGS AND PUPPIES 12 WEEKS OF AGE AND OLDER.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

1. Buckle collar around dog's/puppy's neck. Collar should fit snug enough to prevent placement of the dog's/puppy's leg or lower jaw through the collar, yet loose enough to be rotated about the neck. **2.** Leave 1-2 extra inches on the length of the collar for adjustment, cut off excess, and dispose of in trash. **3.** Check periodically and adjust fit if necessary, especially when puppies are rapidly growing. **4.** Replace in 5 months or sooner if needed. Collar does not need replacement if dog/puppy becomes temporarily wet.

To minimize flea infestation use appropriate ADAMS™ Plus flea and tick products to: **1.** Treat the indoor environment, especially where pets spend time resting. **2.** Treat the outdoor environment when necessary. **3.** Only use EPA-registered products that are labeled for these uses.

STORAGE and DISPOSAL: Storage: Store in original unopened package away from children. **Disposal:** Do not reuse container or collar. Dispose of in trash. Collar should be left in place once fitted on dog/puppy. If collar is temporarily removed, wrap in paper or plastic; do not allow children to have access to collar.

WHAT TO EXPECT:

For Fleas and Ticks:

1. This ADAMS™ Plus Flea & Tick Collar for Small Dogs is designed to kill adult fleas, ticks, and flea eggs over the entire pet. **2.** This collar will provide continuous protection against fleas, ticks, and flea eggs for up to 5 months. **3.** Dead ticks will fall off or may be easily removed. **4.** Replace collar every 5 months or sooner if needed.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS-

CAUTION: Do not get dust or collar in mouth. Harmful if swallowed or absorbed through skin. Do not get dust in eyes, will cause temporary constriction of the pupils*. The dust released by this collar is a cholinesterase inhibitor. Avoid contact with skin, eyes or clothing. **NOTICE: This product contains propoxur, a chemical known to the State of California to cause cancer.**

Humans: Do not open package until ready to use. Do not allow children to play with this collar. Pesticidal dust will form on this collar during storage. Wash hands thoroughly with soap and water after handling collar.

Animals: Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing dogs. Do not use on puppies under 12 weeks of age. Do not use this product on dogs/puppies simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting** drugs or pesticides (e.g. carbaryl or tetrachlorvinphos, etc.). However, flea and tick collars may be immediately replaced. When collar is first worn, observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, remove collar and bathe your dog/puppy with mild soap and rinse with large amounts of water.

FIRST AID

Call a poison control center or doctor immediately for treatment advice.

If swallowed

Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

If in eyes

Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IN CASE OF EMERGENCY, CALL 1-800-234-2269 OR A POISON CONTROL CENTER. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN/VETERINARIAN: Contains an N-methyl carbamate that inhibits cholinesterase.
*Atropine is antidotal only if symptoms of cholinesterase inhibition are present.

To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions.

Satisfaction guaranteed or your money back. Call 1-800-234-2269 or visit our Web site: adamsfleacontrol.com for additional product use information.

**For information on cholinesterase inhibition please call our 800 number or consult your veterinarian.

EPA Reg. No. 2724-493-270 EPA Est. No. 2724-TX-1

CENTRAL Garden & Pet

Manufactured for: **FARNAM PET PRODUCTS**, A Division of Farnam Companies, Inc., P.O. Box 34820, Phoenix, AZ 85067-4820

©2009 Farnam Companies, Inc.

Made in USA

Adams is a licensed trademark and Farnam and the Farnam logo are trademarks of Farnam Companies, Inc.

Central Garden & Pet and its logo are trademarks of Central Garden & Pet Company.

Net Weight:	Product Code:	RM#:	
.69 OZ (19.8 g)	3006024	300505807	09-1190

NAC No.: 10002301