

PFIZER INC.

PFIZER ANIMAL HEALTH

235 E. 42ND ST., NEW YORK, NY, 10017

Telephone: 269-833-4000

Customer Service: 800-733-5500 and 800-793-0596

Veterinary Medical Investigations & Product Support: 800-366-5288

Technical Services (USA): 800-366-5288

Website: <http://pfizerah.com>

NEMEX™ TABS

Pfizer Animal Health

*(pyrantel pamoate)*

### **Canine Anthelmintic Tablets**

*For Small Dogs and Puppies*

**Active Ingredient:** Each tablet contains 22.7 mg of pyrantel base as pyrantel pamoate.

*100 tablets*

*For Large Dogs*

**Active Ingredient:** Each tablet contains 113.5 mg of pyrantel base as pyrantel pamoate.

*50 tablets*

*For veterinary use only*

NADA #100-237, Approved by FDA

For removal of ascarids (*Toxocara canis*; *Toxascaris leonina*) and hookworms (*Ancllyostoma caninum*; *Uncinaria stenocephala*) in dogs and puppies. To prevent reinfection of *T. canis* in puppies and adult dogs and in lactating bitches after whelping.

**Directions:**

*For Small Dogs and Puppies:* For the removal of large roundworms (ascarids) and hookworms, give 1 tablet for each 10 lb of body weight (Dosage is designed to provide at least 2.27 mg per lb of body weight for dogs weighing over 5 lb and at least 4.54 mg per lb of body weight for dogs weighing less than 5 lb). For dogs weighing more than 10 lb, tablets may be broken in half to provide 1/2 tablet for each additional 5 lb of body weight. The presence of these parasites should be confirmed by laboratory fecal examination. Do not withhold food from dog prior to or after treatment.

Place tablet directly into back of mouth or conceal tablet in a small amount of food. A follow-up fecal examination should be conducted 2-4 weeks after first treatment to determine the need for retreatment.

Because anthelmintics cannot be relied upon to prevent reinfection or to remove larvae not present in the intestinal tract at the time of initial treatment, for maximum control, it is recommended that puppies be treated at 2, 3, 4, 6, 8 and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs should be routinely treated at monthly intervals to protect against environmental *T. canis* reinfection. Retreatment of adult dogs may be necessary at monthly intervals as determined by laboratory fecal examinations or in animals kept in known contaminated quarters. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

*For Large Dogs:* For the removal of large roundworms (ascarids) and hookworms, give 1 tablet for each 50 lb of body weight. Tablets may be broken in half to provide 1/2 tablet for 25 lb of body weight. The presence of these parasites should be confirmed by laboratory fecal examination. Do not withhold food from dog prior to or after treatment.

Place tablet directly into back of mouth or conceal tablet in a small amount of food. A follow-up fecal examination should be conducted 2-4 weeks after first treatment to determine the need for retreatment.

Because anthelmintics cannot be relied upon to prevent reinfection or to remove larvae not present in the intestinal tract at the time of initial treatment, for maximum control, it is recommended that puppies be treated at 2, 3, 4, 6, 8 and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs should be routinely treated at monthly intervals to protect against environmental *T. canis* reinfection. Retreatment of adult dogs may be necessary at monthly intervals as determined by laboratory fecal examinations or in animals kept in known contaminated quarters. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

**Caution:** If dog looks or acts sick, do not treat with this product.

**Caution:** Keep out of reach of children. For use only as directed on label.

**Restricted Drug (California)-Use Only as Directed**

**Not for Human Use**

*Distributed by:* **Pfizer Animal Health**, Exton, PA 19341, USA, Div. of Pfizer Inc. NY, NY 10017

015 01-4166-35-3

014 01-4167-35-3

Made in USA

**NAC No.:** 36901201

