

PFIZER INC.

PFIZER ANIMAL HEALTH

235 E. 42ND ST., NEW YORK, NY, 10017

Telephone: 269-833-4000

Customer Service: 800-733-5500 and 800-793-0596

Veterinary Medical Investigations & Product Support: 800-366-5288

Technical Services (USA): 800-366-5288

Website: <http://pfizerah.com>

STRONGID® T




Pfizer Animal Health

(pyrantel pamoate)

Equine Anthelmintic Suspension

Strongid T is a suspension of pyrantel pamoate in a palatable caramel-flavored vehicle. Each mL contains 50 mg of pyrantel base as pyrantel pamoate.

Pyrantel pamoate is a compound belonging to a family classified chemically as tetrahydropyrimidines. It is a yellow, water-insoluble crystalline salt of the tetrahydropyrimidine base and pamoic acid containing 34.7% base activity. The chemical structure and name are given below:



(E)-1,4,5,6-Tetrahydro-1-methyl-2-[2-(2-thienyl) vinyl] pyrimidine 4,4' methylenebis [3- hydroxy-2-naphthoate] (1:1)

INDICATIONS FOR USE: For the removal and control of mature infections of large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*); pinworms (*Oxyuris equi*); large roundworms (*Parascaris equorum*); and small strongyles in horses and ponies.

DOSAGE AND TREATMENT: Administer 3 mg pyrantel base per lb of body weight (6 mL Strongid T per 100 lb of body weight). For maximum control of parasitism, it is recommended that foals (2-8 months of age) be dosed every 4 weeks. To minimize potential hazard that the mare may pose to the foal, she should be treated 1 month prior to anticipated foaling date followed by retreatment 10 days to 2 weeks after birth of foal. Horses over 8 months of age should be routinely dosed every 6 weeks.

DIRECTIONS FOR USE: Strongid T may be administered by means of a stomach tube, dose syringe or by mixing into the feed.

Stomach Tube: Measure the appropriate dosage of Strongid T and mix in the desired quantity of water. Protect drench from direct sunlight and administer to the animal immediately following mixing. Do not attempt to store diluted suspension.

Strongid T is inactive against the common horse bot (*Gasterophilus* sp.). However, Strongid T may be administered concurrently with carbon disulfide observing the usual precautions with carbon disulfide.

Dose Syringe: Draw the appropriate dosage of Strongid T into a dose syringe and administer to the animal. Do not expose Strongid T to direct sunlight.

Feed: Mix the appropriate dosage of Strongid T in the normal grain ration. Fasting of animals prior to or following treatment is not required.

CAUTION: THIS PRODUCT IS A SUSPENSION AND AS SUCH WILL SEPARATE. TO INSURE UNIFORM RESUSPENSION AND TO ACHIEVE PROPER DOSAGE, IT IS EXTREMELY IMPORTANT THAT THE PRODUCT BE SHAKEN AND STIRRED THOROUGHLY BEFORE EVERY USE.

EFFICACY: Critical (worm-count) studies in horses demonstrated that Strongid T administered at the recommended dosage was efficacious against mature infections of *Strongylus vulgaris* (>90%), *S. edentatus* (69%), *S. equinus* (>90%), *Oxyuris equi* (81%), *Parascaris equorum* (>90%), and small strongyles (>90%).

SAFETY: Strongid T is well tolerated by horses and ponies of all ages. No adverse drug response was observed when dose rates up to 60 mg pyrantel base per lb of body weight were administered by stomach tube nor when 3 mg base per lb was given by intratracheal injection. The reproductive performance of pregnant mares and stud horses dosed with Strongid T has not been affected.

▶ **WARNINGS:** Not for horses or ponies intended for food. Keep out of the reach of children. ◀

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

It is recommended that severely debilitated animals not be treated with this preparation.

RECOMMENDED STORAGE: Store below 30°C (86°F).

NADA #91-739, Approved by FDA

Distributed by: **Pfizer Animal Health**, Exton, PA 19341, USA, Div. of Pfizer Inc, NY, NY 10017

004

75-7972-00

Net Contents:	
1 quart (946 mL)	85-7972-00

NAC No.: 36901432